

# **HERPES SIMPLEX SURVEILLANCE PROTOCOL FOR ONTARIO HOSPITALS**

Developed by the Ontario Hospital Association and  
the Ontario Medical Association  
Joint Communicable Diseases Surveillance Protocols Committee

Approved by  
The OHA and the OMA Board of Directors  
The Ministry of Health and Long-Term Care –  
The Minister of Health and Long-Term Care

Published and Distributed by the Ontario Hospital Association  
Published September 1991  
Last Reviewed and Revised November 2018

# Herpes Simplex Surveillance Protocol for Ontario Hospitals

Published September 1991  
Last Reviewed and Revised November 2018

This protocol was developed jointly by the Ontario Hospital Association and the Ontario Medical Association to meet the requirements of the *Public Hospitals Act 1990*, Revised Statutes of Ontario, Regulation 965. This regulation requires each hospital to have by-laws that establish and provide for the operation of a health surveillance program including a communicable disease surveillance program in respect of all persons carrying on activities in the hospital. The communicable disease program is to include the tests and examinations set out in any applicable communicable disease surveillance protocol. The regulation states that the communicable disease surveillance protocols that hospitals must adopt are those "published jointly by the Ontario Hospital Association (OHA) and the Ontario Medical Association (OMA) and approved by the Minister (of Health and Long-Term Care)."

This Protocol has been reviewed since the previous version; changes have been highlighted in yellow for easy identification. Protocols are reviewed on a regular basis, every two years or as required.

The protocol reflects clinical knowledge, current data and experience, and a desire to ensure maximum cost effectiveness of programs, while protecting health care workers and patients. It is intended as a minimum standard that is practical to apply in most Ontario hospital settings. It does not preclude hospitals from adopting additional strategies that may be indicated by local conditions.

# Members of the Joint OHA/OMA Communicable Disease Surveillance Protocols Committee

## Representing the Ontario Hospital Association

Dr. Kathryn Suh (Co-chair)  
Medical Director, Infection Prevention and Control Program  
The Ottawa Hospital, Ottawa

Kathleen Poole, MScN, COHN(C) CIC  
Infection Control Practitioner,  
Providence Care, Kingston

Sandra Callery, RN MHSc CIC  
Director, Infection Prevention and Control

## Ontario Hospital Association

Laurie Cabanas  
Acting Director, Policy

Amanda Martens  
Policy Advisor

## Representing the Ontario Medical Association

Dr. Maureen Cividino (Co-chair)  
IPAC Physician, Public Health Ontario  
Occupational Health Physician  
St. Joseph's Healthcare, Hamilton

Dr. Irene Armstrong  
Associate Medical Officer of Health  
Communicable Disease Control  
Toronto Public Health, Toronto

## Ontario Medical Association

Katherine Patterson  
Senior Advisor, Health Policy and Promotion  
Ontario Medical Association

## Representing the Ministry of Health and Long-Term Care

Melissa Helferty, MIPH  
Manager, Infectious Disease Policy & Programs  
Health Protection and Surveillance Policy and Programs Branch

## EX-OFFICIO

Dr. Nikhil Rajaram  
Senior Medical Consultant, Occupational  
Medicine Unit  
Occupational Health and Safety Branch  
Ministry of Labour

Henrietta Van hulle, BN, MHSM, COHN(c), CRSP,  
CDMP  
Vice President, Client Outreach  
Public Services Health and Safety Association

# Rationale for Herpes Simplex Surveillance Protocol

Herpes simplex virus (HSV) is an extremely common cause of infection worldwide, with 50-90% of adults having antibodies.<sup>1</sup> Direct contact with lesions or infected secretions is the primary mode of transmission of HSV.

Primary infection with HSV infection results from the first exposure to the virus. Orolabial infection may be mild or inapparent, or result in overt disease with fever, malaise lasting a week or more and vesicular lesions in the mouth and/or pharynx. After primary infection, the virus becomes latent, and may reactivate as localized blisters, commonly referred to as “cold sores” or “fever blisters”, which usually appear at the border between the mucous membranes of the mouth and the skin. Genital infections, skin lesions, eye infections, generalized systemic infections and central nervous system infections may also occur with primary infection and reactivation.

Primary or recurrent lesions, and secretions such as saliva and genital secretions, contain the virus. Virus may be present in saliva or genital secretions in the absence of symptomatic or clinically apparent lesions.<sup>2</sup>

Health care workers (HCWs) are at risk of acquiring HSV infection if exposed hands contact infected secretions or mucous membranes/skin of an asymptomatic or symptomatic patients shedding the virus.<sup>3,4</sup> Infections acquired by this route typically result in herpetic whitlow, painful recurring vesicular lesions on the nail or finger area, although other herpes infections may also occur.<sup>5</sup> Dentists and dental teams,<sup>6-8</sup> anaesthesiologists,<sup>9</sup> respiratory therapists, obstetricians, and critical care nurses<sup>5,7,10</sup> are at particular risk. Routine Practices, including wearing gloves,<sup>11,12</sup> reduce this risk.

Patients are also at risk of acquiring HSV from infected HCWs. For some high risk patients (e.g. newborns, patients with extensive skin damage such as burns and chronic eczema, patients who are immunocompromised) infection may result in severe, life-threatening, systemic disease. Virus can be transferred from the oral area to the hands of the HCW, and then transmitted by HCW hands on contact with the patient. Although rare, transmission from HCWs to newborns has been documented.<sup>13-15</sup>

Because HSV may be present in saliva before or without the appearance of lesions, continuing education must emphasize the mode of transmission, the importance of hand hygiene and use of Routine Practices when in direct patient contact to prevent transmission both from patients to HCWs and from HCWs to patients.

**This protocol is only one component of an infection prevention and control program; HCWs must consistently adhere to Routine Practices.**

# Herpes Simplex Surveillance Protocol for Ontario Hospitals

Developed by  
the Ontario Hospital Association and the Ontario Medical Association  
Published September 1991  
Last Reviewed and Revised November 2018

## I. Purpose

The purpose of this protocol is to provide direction to hospitals to prevent the transmission of herpes simplex virus among health care workers (HCWs) and patients.

## II. Applicability

This protocol applies to **all persons carrying on activities in the hospital**, including but not limited to employees, physicians, nurses, contract workers, students, post-graduate medical trainees, researchers and volunteers. The term HCW is used in this protocol to describe these individuals. This protocol does not apply to patients or residents of the facility or to visitors.

When training students or hiring contract workers, the hospital must inform the school/supplying agency that the school/agency is responsible for ensuring that their student/contractors are managed according to this protocol.

**This protocol is for the use of the Occupational Health Service (OHS) in hospitals. It is expected that OHS collaborate with Infection Prevention and Control (IPAC) and other departments, as applicable.**

## III. Pre-placement

Screening for HSV in persons carrying on activities in the hospital is neither required nor recommended. HCWs must be informed of the requirement to notify the OHS of acute primary herpes simplex infection, and recurrent orofacial infection or herpetic whitlow.

## IV. Continuing Surveillance

No routine screening for HSV in persons carrying on activities in the hospital is needed or recommended.

## V. Exposure

Exposure has occurred if unprotected hands have come into contact with infected secretions or mucous membranes/skin of asymptomatic or symptomatic patients shedding the virus.<sup>3,4</sup>

## VI. Acute Disease

HSV infections are evident as:

- oral infections (inside the mouth),
- orofacial infections (inside the mouth and on the outside of the lips, face),
- herpetic whitlow (herpes simplex infection of the thumb or fingers),
- genital infections, and
- herpes corporis (herpes simplex infection on parts of the body other than hands, face or genitals).

Persons carrying on activities in the hospital who have direct patient contact and who develop acute infections of the first three types below (oral, orofacial, herpetic whitlow) have a responsibility to inform the OHS as soon as they notice symptoms. Work restrictions or modifications vary with the type of herpes simplex infection and the type of patients with whom the person has contact. HCWs excluded from direct patient contact because of HSV infection may safely be assigned to duties that involve no patient contact.

### Work Restrictions

Acute Primary Oropharyngeal Infection:

- No patient contact until symptoms have resolved.

Oral/Orofacial Infection:

- HCWs working with high-risk patients (see Glossary) may continue to work as long as they **maintain meticulous hand hygiene, and cover the lesions (e.g., wear a surgical mask or dressing)** to discourage hand-to-lesion contact. The HCW should **wear gloves for direct hands-on contact with high risk patients**. Perform hand hygiene after removing gloves, as per Routine Practices.
- HCWs who do not work with high-risk patients may continue to work with no special precautions, as long as they maintain meticulous hand hygiene.

Herpetic Whitlow:

- HCWs with HSV infection of the fingers must be restricted from all direct patient contact until lesions are healed.<sup>11</sup> There is no evidence

that wearing gloves will provide adequate protection for the patient, and prolonged wearing of gloves would likely aggravate the existing infection.

#### Genital Herpes / Herpes Corporis

- HCWs with genital HSV infection or herpes corporis do not have to inform OHS of these conditions. There should be no risk to patients as long as good hygiene is maintained, including meticulous hand hygiene, and lesions are covered.

## Work Restrictions Table

Note direct patient care refers to skin-to-skin contact in patient care activities that require direct, personal “hands-on” care (e.g., bathing, washing, turning patient, changing clothes, continence care, dressing changes, care of open wounds/lesions, toileting), (OAHPP, 2012).

Primary HSV Infection *	Location of Vesicular Lesions	Work Restrictions/Practices
Acute primary oropharyngeal infection	Mouth, pharynx, oral cavity +/- tonsils	No patient contact until symptoms resolved
Reactivation HSV Infection*	Location of Vesicular Lesions	Work Restrictions/Practices
Oral/Orofacial	Inside mouth, outside of lips, face	<p><b>Patients not high risk:</b> No restrictions Maintain meticulous hand hygiene</p> <p><b>High risk patients</b> (newborns, burns, extensive skin damage): Cover lesions (mask/dressing) for direct patient care to avoid hand to lesion contact Wear gloves for all direct hands-on contact</p>
Herpetic whitlow	Thumb or fingers	<p>No direct patient care until lesions are healed</p> <p><b>No evidence that gloves protect against transmission of infection</b></p>
Herpes Corporis	Areas on the body other than the hands, face or genitals	No work restrictions or need to report to OHS

\*Genital herpes: No work restrictions or need to report to OHS



## **VII. Reporting**

Herpes simplex infections are not reportable to the Medical Officer of Health.

In accordance with the Occupational Health and Safety Act and its regulations, an employer must provide written notice within 4 days of being advised that a worker has an occupational illness, including an occupationally-acquired infection, and/or a Workplace Safety and Insurance Board (WSIB) claim has been filed by or on behalf of the worker with respect to an occupational illness, including an occupational infection, to the:

- Ministry of Labour,
- Joint Health and Safety Committee (or health and safety representative), and
- trade union, if any.

Occupationally-acquired infections and illnesses are reportable to the WSIB.

## **VIII. Outbreak**

Not applicable in this protocol.

## References

1. Xu F, Sternberg MR, Kottiri BJ, et al. Trends in herpes simplex virus type 1 and 2 seroprevalence in the United States. *JAMA* 2006;296:964-73.
2. Miller CS, Danaher RJ. Asymptomatic shedding of herpes simplex virus (HSV) in the oral cavity. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;105:43-50.
3. Chayavichitsilp P, Buckwalter JV, Krakowski AC, Friedlander SF. Herpes simplex. *Pediatrics in Review* 2009;30:119-29.
4. Whitley RJ. Herpes simplex encephalitis: adolescents and adults. *Antiviral Res* 2006;71:, 141-8.
5. Perl TM, Haugen TH, Pfaller MA, et al. Transmission of herpes simplex virus 1 infection in an intensive care unit. *Ann Int Med* 1992;117:584-6.
6. Lewis MA. Herpes simplex virus: an occupational hazard in dentistry. *Int Dent J* 2004;54:103-11.
7. Rowe NH, Heine CS, Kowalski CJ. Herpetic whitlow: an occupational disease of practicing dentists. *J Am Dent Assoc* 1982;105:471-3.
8. Browning WD, McCarthy JP. A case series: Herpes simplex virus as an occupational hazard. *J Esthet Restor Dent* 2012;24:61-6.
9. Orkin FK. Herpetic whitlow – occupational hazard to the anesthesiologist. *Anesthesiology* 1970;33:671-3.
10. Adams G, Stover BH, Keenlyside RA et al. Nosocomial herpetic infections in a pediatric intensive care unit. *Am J Epidemiol* 1981;113:126-32.
11. Health Canada. Prevention and control of occupational infections in health care. *Can Comm Dis Rep* 2002;28(S1).
12. Ontario Agency for Health Protection and Promotion, Provincial Infectious Diseases Advisory Committee. Routine practices and additional precautions in all health care settings. Toronto, ON: Queen's Printer for Ontario. November 2012.
13. Sakaoka H, Saheki Y, Uzuki K, et al. Two outbreaks of herpes simplex virus type 1 nosocomial infection among newborns. *J Clin Microbiol* 1986; 24:36-40.
14. Hammerberg O, Watts J, Chernesky M, Luchsinger I, Rawls W. An outbreak of herpes simplex virus type 1 in an intensive care nursery. *Pediatr Infect Dis J* 1983;2:290-4.

15. Linnemann CC Jr, Buchman TG, Light IJ, Ballard JL. Transmission of herpes simplex virus type 1 in a nursery for the newborn: identification of viral isolates by DNA fingerprinting. *Lancet* 1978;1(8071):964-6.