

## **GET THE MAXIMUM USE**

from each mask that safety allows to reduce overuse and reduce the risk of running out. This is the most important piece of teamwork we can do.

#### REGULAR PATIENT CONTACT IN A COVID-DEDICATED AREA

- Wear mask-with-visor for 4 hours.
- Once on, leave it in place. Don't adjust or touch.
- Remove/discard on break or lunch.

## NOT IN A PATIENT CARE AREA

- If your job can be done at home, it must be done at home.
- If you must be in the hospital, keep
   2 meters distance between you and all other people.

# WORK IN PATIENT CARE AREAS BUT NOT REGULARLY INTERACTING WITH PATIENTS ON DROPLET PRECAUTIONS

- Wear your procedure mask as long as possible, and up to 4 hours.
- Once on, leave it in place. Don't adjust or touch.
- Remove it to eat and place it in a bag or clean surface and then use it again if less than 4 hours.
- When in contact with patients on droplet precautions, add face shield. (If no face shield, change to a maskwith-visor OR add goggles).

### SPECIAL INSTRUCTIONS FOR DI AND STAFF WHO ROVE FROM UNIT TO UNIT LIKE:

- Dietary
- RTs
- Other Allied Health

See full protocol for details

MASKS THAT BECOME SOILED CAN BE REPLACED WHEN NECESSARY

**DON'T SKIMP and DON'T WASTE**